

N65-29492

(ACCESSION NUMBER)

(THRU)

(PAGES)

(CODE)

(NASA CR OR TMX OR AD NUMBER)

(CATEGORY)

GPO PRICE \$

CFSTI PRICE(S) \$

UNPUBLISHED

The Fourier Series of Gegenbauer's Function

Hard copy (HC) 1.00

B. C. Carlson and W. H. Greiman

Microfiche (MF) 50

Iowa State University, Ames, Iowa

ff 653 July 65

1. Introduction. If N is a positive integer, the Gegenbauer polynomial C_N^ν is known to have the representation [3, vol. 2, p. 175]

$$(1.1) \quad C_N^\nu(\cos \theta) = \sum_{m=0}^N \frac{(\nu)_m (\nu)_{N-m}}{m! (N-m)!} \cos(N-2m)\theta,$$

where $(\nu)_m = \Gamma(\nu+m)/\Gamma(\nu)$. The Fourier series of Gegenbauer's function $C_\alpha^\nu(\cos \theta)$ with general (possibly complex) α does not appear to have been given previously, even in the special case of Legendre's function $P_\alpha = C_\alpha^{\frac{1}{2}}$.

We shall find that

$$(1.2) \quad C_\alpha^\nu(\cos \theta) = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n \cos n\theta, \quad (\operatorname{Re} \nu < 1),$$

$$A_n = \left\{ 1 + \frac{\sin \pi(\nu + \alpha + n)}{\sin \pi \nu} \right\} \frac{\Gamma(\nu + \frac{\alpha+n}{2}) \Gamma(\nu + \frac{\alpha-n}{2})}{[\Gamma(\nu)]^2 \Gamma(1 + \frac{\alpha+n}{2}) \Gamma(1 + \frac{\alpha-n}{2})}.$$

If $\operatorname{Re} \nu \geq 1$ the Fourier coefficients do not exist for general α because $C_\alpha^\nu(\cos \theta)$ is not integrable over an interval containing the point $\theta = \pi$.

If $\operatorname{Re} \nu < 1$ and α is a positive integer N , the first factor of A_n vanishes if $N+n$ is odd; since $A_{-n} = A_n$, (1.2) then reduces to (1.1).

We remark that Gegenbauer's function (multiplied by a constant to

¹ This work was supported by the Ames Laboratory of the U. S. Atomic Energy Commission and by the National Aeronautics and Space Administration under Grant NsG-293 to Iowa State University.

give it the value unity at $\theta = 0$) has a nicely symmetrical expression in the notation of the hypergeometric R function [1]:

$$(1.3) \quad \frac{\Gamma(2\nu) \Gamma(\alpha + 1)}{\Gamma(2\nu + \alpha)} C_{\alpha}^{\nu}(\cos \theta) = R(-\alpha; \nu, \nu; e^{i\theta}, e^{-i\theta}) \\ = 2^{\nu-\frac{1}{2}} \Gamma(\nu + \frac{1}{2}) (\sin \theta)^{\frac{1}{2}-\nu} P_{\alpha+\nu-\frac{1}{2}}^{\frac{1}{2}-\nu}(\cos \theta),$$

where P is an associated Legendre function. An important special case is

$$(1.4) \quad C_{\alpha}^{\frac{1}{2}}(\cos \theta) = R(-\alpha; \frac{1}{2}, \frac{1}{2}; e^{i\theta}, e^{-i\theta}) = P_{\alpha}(\cos \theta).$$

2. The Fourier coefficients. Gegenbauer's function is defined [3, vol. 1, p. 178] by

$$(2.1) \quad \frac{\Gamma(2\nu) \Gamma(\alpha + 1)}{\Gamma(2\nu + \alpha)} C_{\alpha}^{\nu}(\cos \theta) = {}_2F_1(-\alpha, 2\nu + \alpha; \nu + \frac{1}{2}; \sin^2 \frac{\theta}{2}) \\ = \sum_{m=0}^{\infty} \frac{(-\alpha)_m (2\nu + \alpha)_m}{(\nu + \frac{1}{2})_m m!} \sin^{2m}(\theta/2).$$

We assume for the present that $\operatorname{Re} \nu < \frac{1}{2}$, so that the hypergeometric series converges absolutely for $\theta = \pi$ [5, p. 25] and hence uniformly over the interval $(0, \pi)$. The Fourier coefficient

$$(2.2) \quad A_n = \frac{2}{\pi} \int_0^{\pi} C_{\alpha}^{\nu}(\cos \theta) \cos n\theta \, d\theta$$

can then be found by integrating term by term. From the elementary formula

$$(2.3) \quad \frac{2}{\pi} \int_0^{\pi} \sin^{2m}(\theta/2) \cos n\theta \, d\theta = (-1)^n 2^{1-2m} \binom{2m}{m+n},$$

we get

$$\begin{aligned}
 A_n &= \frac{(-1)^n 2 \Gamma(2\nu + \alpha)}{\Gamma(2\nu) \Gamma(\alpha + 1)} \sum_{m=n}^{\infty} \frac{(-\alpha)_m (2\nu + \alpha)_m (\frac{1}{2})_m}{(\nu + \frac{1}{2})_m (m+n)! (m-n)!} \\
 (2.4) \quad &= \frac{(-1)^n 2^{1-2n} \Gamma(2\nu + \alpha + n) (-\alpha)_n}{\Gamma(2\nu) \Gamma(\alpha + 1) (\nu + \frac{1}{2})_n n!} \sum_{k=0}^{\infty} \frac{(n - \alpha)_k (2\nu + \alpha + n)_k (\frac{1}{2} + n)_k}{(\nu + \frac{1}{2} + n)_k (1 + 2n)_k k!}.
 \end{aligned}$$

The last series, obtained from the preceding one by putting $m = n + k$, is a ${}_3F_2$ series with unit argument. If $\operatorname{Re} \nu < 1$ it converges and can be summed by Watson's theorem [3, vol. 1, p. 189]:

$$\begin{aligned}
 (2.5) \quad {}_3F_2(n - \alpha, 2\nu + \alpha + n, \frac{1}{2} + n; \nu + \frac{1}{2} + n, 1 + 2n; 1) \\
 = \frac{\pi^{\frac{1}{2}} n! \Gamma(\nu + \frac{1}{2} + n) \Gamma(1 - \nu)}{\Gamma(\frac{1 + n - \alpha}{2}) \Gamma(\frac{1 + n + 2\nu + \alpha}{2}) \Gamma(1 + \frac{n + \alpha}{2}) \Gamma(1 + \frac{n - 2\nu - \alpha}{2})}.
 \end{aligned}$$

Substitution in (2.4) gives an expression for A_n that can be simplified by applying several times the duplication formula for the gamma function and the relation $\Gamma(z) \Gamma(1 - z) = \pi \csc \pi z$. The result is

$$(2.6) \quad A_n = \frac{(-1)^n \sin \pi \alpha \sin \pi(\nu + \frac{\alpha - n}{2}) \Gamma(\nu + \frac{\alpha + n}{2}) \Gamma(\nu + \frac{\alpha - n}{2})}{\sin \pi \nu \sin \pi(\frac{\alpha - n}{2}) [\Gamma(\nu)]^2 \Gamma(1 + \frac{\alpha + n}{2}) \Gamma(1 + \frac{\alpha - n}{2})}.$$

Elementary rearrangement of the sine functions now leads to (1.2).

If $\operatorname{Re} \nu \geq \frac{1}{2}$ the series (2.1) no longer converges uniformly over $(0, \pi)$ if it does not terminate. However, the analytic continuation of Gauss' hypergeometric function [5, p. 291] shows that $C_{\alpha}^{\nu}(\cos \theta)$ is then of the order of $(\cos \frac{1}{2}\theta)^{1-2\nu}$ as $\theta \rightarrow \pi$ (except that the singularity is logarithmic if $\nu = \frac{1}{2}$).

Provided that $\operatorname{Re} \nu < 1$, the function is integrable over $(0, \pi)$; furthermore, it satisfies conditions [5, p. 164] sufficient to ensure that its Fourier series converges (except when θ is an odd multiple of π) and represents the function.

To show that the Fourier coefficients are still given by (1.2) if $\frac{1}{2} \leq \operatorname{Re} \nu < 1$, one can either use analytic continuation in ν or justify directly the term-by-term integration of (2.1). The second method is the easier if one uses the following theorem [4, p. 45]: If $\sum u_m(\theta)$ converges uniformly over $(0, \pi - \epsilon)$ for every (small) positive ϵ , and if $\sum \int_0^\pi |u_m(\theta)| d\theta$ converges, then $\sum u_m(\theta)$ may be integrated term by term over $(0, \pi)$.

The first assumption of the theorem is plainly satisfied by

$$u_m(\theta) = \frac{(-\alpha)_m (2\nu + \alpha)_m}{(\nu + \frac{1}{2})_m m!} \sin^{2m}(\theta/2) \cos n\theta .$$

Moreover, we have

$$\begin{aligned} \sum \int_0^\pi |u_m(\theta)| d\theta &\leq \sum \left| \frac{(-\alpha)_m (2\nu + \alpha)_m}{(\nu + \frac{1}{2})_m m!} \right| \int_0^\pi \sin^{2m}(\theta/2) d\theta \\ &= \pi \sum \left| \frac{(-\alpha)_m (2\nu + \alpha)_m (\frac{1}{2})_m}{(\nu + \frac{1}{2})_m m! m!} \right| . \end{aligned}$$

The last series converges if $\operatorname{Re} \nu < 1$, and the proof of (1.2) is now complete.

Eq. (1.3) follows from (2.1) by use of the relation [1, Eq. (2.5)]

$$(2.7) \quad {}_2F_1(a, b; c; x) = R(a; b, c - b; 1 - x, 1)$$

and the quadratic transformation [2, Eq. (5.1)]

$$(2.8) \quad R(a; b, b; x^2, y^2) = R(a; 2b - a, a - b + \frac{1}{2}; (x + y)^2/4, xy) .$$

References

1. B. C. Carlson, Lauricella's hypergeometric function F_D , Journal of Mathematical Analysis and Applications, vol. 7 (1963), pp. 452-470.
2. B. C. Carlson, Normal elliptic integrals of the first and second kinds, Duke Mathematical Journal, September, 1964.
3. A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, Higher Transcendental Functions, New York, 1953.
4. E. C. Titchmarsh, The Theory of Functions, 2nd ed., Oxford, 1939.
5. E. T. Whittaker and G. N. Watson, Modern Analysis, 4th ed., Cambridge, 1927.